

The π –Gluon Exchange Interaction Between Constituent Quarks

C. Helminen and D.O. Riska

Department of Physics, University of Helsinki, 00014 Finland

Abstract

The interaction mediated by irreducible pion and gluon exchange between constituent quarks is calculated and shown to have a strong tensor component, which tends to cancel the pion exchange tensor interaction between quarks. Its spin-spin component is somewhat weaker than the pion exchange spin-spin interaction, while its central and spin-orbit components are small in comparison to the corresponding single gluon exchange interactions. The combination of the π –gluon exchange interaction with the single pion exchange interaction and a weak gluon exchange interaction between constituent quarks has the qualitative features required for understanding the hyperfine splittings of the spectra of the nucleon and the Δ resonances.

1. Introduction

Both the form and the dynamical origin of the effective interaction between the constituent quarks, which form the baryons, remain largely open issues. The small spin-orbit splittings between the lowest negative parity states in the baryon spectrum suggest that both the spin-orbit and tensor components of the effective quark-quark interaction should be weak. This rules out a strong quark-gluon coupling, as does the presence of low lying positive parity states below the lowest negative parity states in all those sectors of the baryon spectrum without flavor singlet states. The latter feature is most readily explained if the main component of the hyperfine interaction between the quarks is an attractive flavor dependent spin-spin interaction [1].

Such an interaction is an integral component of the Goldstone boson (i.e. π, K, η) exchange interaction between quarks, and as long as the associated tensor component is dropped, it is indeed possible to explain the spectrum of the nucleon and the strange hyperons up to the small spin-orbit splittings with that interaction in combination with a linear confining interaction with conventional strength [2]. This however leaves the tensor component of the Goldstone boson exchange interaction to be explained away, somewhat as the large spin-orbit interaction of the gluon exchange interaction has to be explained away in attempts to describe the baryon spectrum in terms of gluon exchange alone [3].

We here point out that if the Goldstone boson exchange interaction between quarks is complemented with a fairly weak gluon exchange interaction along with the associated irreducible π -gluon exchange interaction the problem of the tensor component of the former is ameliorated if not eliminated. This is because the tensor component of the π -gluon exchange interaction is of the same order of magnitude as the pion exchange interaction, but has the opposite sign. As a result the net tensor interaction is very weak. This provides an alternative to balancing the pion exchange tensor interaction at short range by a corresponding vector meson exchange interaction of opposite sign [1].

The π -gluon exchange interaction has an attractive spin-spin component, which adds to, but is somewhat weaker than, that of single pion exchange at short range. Its detailed behavior at very short range is very sensitive to the high momentum behavior of the pion and gluon exchange interactions. Finally the spin-orbit and central components of the π -gluon exchange interaction turns out to be very weak. The π -gluon exchange interaction thus appears to provide part of the explanation for why the effective interaction between constituent quarks should have the form of an attractive flavor dependent spin-spin interaction and an at most very weak tensor interaction.

Another issue is to what extent the nonperturbative vacuum of QCD supports gluon exchange. Cooled lattice calculations suggests that the quark-gluon coupling should be very weak [4]. The valence-QCD approximation suggests the presence of a residual weak but nonzero gluon exchange interaction between quarks [5]. We shall here assume that the gluons either decouple from the constituent quarks below the confinement scale Λ_{QCD} or the chiral restoration scale $\Lambda_\chi \simeq 1$ GeV or that their coupling freezes at some small value below

these momentum scales. This agrees with recent phenomenological studies of the behavior of the running QCD coupling strength in the infrared limit [6, 7]. In addition we make the conventional assumption that the Goldstone bosons decouple above the chiral restoration scale. The π –gluon exchange loop mechanisms allow combination of short range gluon exchange with long range pion exchange.

For the calculation of the π –gluon exchange interaction we adopt the Blankenbecler-Sugar quasipotential framework, which allows a covariant extraction of the iterated single pion and gluon exchange interactions from the Bethe-Salpeter equation kernel [8]. The resulting interaction is therefore real and almost energy independent. In the calculation we rely largely on the formalism developed in Refs. [9, 10] for a different application.

This paper falls into 7 sections. In section 2 we define the π –gluon exchange interaction. In section 3 we calculate the π –gluon exchange loop diagrams. In section 4 we derive the iterated pion and gluon exchange interaction, which has to be subtracted from the covariant loop diagrams in order to obtain a properly defined irreducible interaction. In section 5 the adiabatic limit of the π –gluon exchange interaction is obtained. In section 6 we give numerical results for the interaction components. Section 7 contains a summarizing discussion.

2. Definition of the π –gluon exchange interaction

The Bethe-Salpeter equation for the invariant quark-quark scattering amplitude M may be written as

$$M = K + KGM, \quad (2.1)$$

where K is the two-quark irreducible interaction kernel and G is the two quark propagator. The conventional assumption, which underlies most quark model phenomenology, is that this may be approximated as a product of free fermion propagators with effective constituent quark masses. The Blankenbecler-Sugar reduction amounts to the separation of this propagator into a quasi-3-dimensional part $\tilde{g}$ and a residual part $G - \tilde{g}$ [8, 11]. This separation reduces the Bethe-Salpeter equation (2.1) to a 3-dimensional quasipotential equation:

$$M = U + U\tilde{g}M, \quad (2.2)$$

with an accompanying defining equation for the quasipotential U :

$$U = K + K(G - \tilde{g})U. \quad (2.3)$$

For the 3-dimensional propagator $\tilde{g}$ we employ the symmetrical choice [12, 13]:

$$\tilde{g}(\vec{k}; W) = 2\pi i \delta(k_0) \frac{\Lambda_+^{(1)}(\vec{k}) \Lambda_+^{(2)}(-\vec{k})}{E^2(\vec{k}) - W^2 - i\epsilon}. \quad (2.4)$$

Here W is the total energy and $\Lambda_+^{(i)}(\vec{k})$ ($i=1,2$) are positive energy projection operators

$$\Lambda_+^{(i)}(\vec{k}) = \frac{\gamma_4^{(i)} E(\vec{k}) - i\vec{\gamma}^{(i)} \cdot \vec{k} + m_q}{2m_q}, \quad (2.5)$$

where m_q is the constituent quark mass.

Consider now the case, where the irreducible interaction kernel K is approximated as the sum of the one-pion- and one-gluon-exchange and the crossed two-pion, two-gluon and pion+gluon exchange interactions:

$$K \simeq K_\pi + K_G + K_{\pi\pi} + K_{GG} + K_{\pi G}. \quad (2.6)$$

The lowest (second) order term in the quasipotential (2.3), which corresponds to the tree approximation is then the sum of one-pion and one-gluon exchange interactions:

$$U^{(2)} = K_\pi + K_G. \quad (2.7)$$

The expressions for these are readily obtained from the pion- and gluon-quark coupling Lagrangians:

$$\begin{aligned} \mathcal{L}_{\pi qq} &= i \frac{f_{\pi qq}}{m_\pi} \bar{\psi} \gamma_\mu \gamma_5 \partial_\mu \vec{\phi} \cdot \vec{\tau} \psi, \\ \mathcal{L}_{Gqq} &= i \sqrt{4\pi\alpha_S} \bar{\psi} \gamma_\mu \vec{G}_\mu \cdot \vec{\lambda}_C \psi. \end{aligned} \quad (2.8)$$

Here $\vec{\lambda}_C$ is the vector of $SU(3)$ color generators and $\vec{G}_\mu$ is the gluon field. For the one-pion and one-gluon exchange interactions between quarks one then obtains the expressions:

$$K_\pi = \frac{f_{\pi qq}^2}{m_\pi^2} \frac{(\gamma^{(1)} \cdot k) \gamma_5^{(1)} (\gamma^{(2)} \cdot k) \gamma_5^{(2)}}{m_\pi^2 + k^2} f(k^2) \vec{\tau}^1 \cdot \vec{\tau}^2, \quad (2.9a)$$

$$K_G = -\frac{8\pi}{3} \alpha_S \frac{\gamma^{(1)} \cdot \gamma^{(2)}}{k^2} g(k^2). \quad (2.9b)$$

Here $f_{\pi qq}$ is the pion-quark coupling constant, and α_S the effective coupling constant for gluons to constituent quarks. The function $f(k^2)$ is a cut-off factor, that should describe the decoupling of the pions from constituent quarks above the chiral restoration scale Λ_χ , so that $f(k^2 = -m_\pi^2) = 1$ and $\lim_{k^2 \rightarrow \infty} f(k^2) = 0$.

We shall here take the cut-off function $f(k^2)$ to have the form

$$f(k^2) = \frac{\Lambda_\chi^2 - m_\pi^2}{\Lambda_\chi^2 + k^2}, \quad (2.10)$$

where Λ_χ is the chiral symmetry restoration scale. The pseudovector pion-quark coupling constant $f_{\pi qq}$ may be related to the πNN pseudovector coupling as $f_{\pi qq} = (3/5)f_{\pi NN} \simeq 0.6$, a value which will be adopted here.

The function $g(k^2)$ on the other hand is a factor introduced to describe the momentum dependence of the quark-gluon coupling constant. For momenta above Λ_{QCD} this factor should contain the inverse logarithmic fall-off factor $1/\log(k^2/\Lambda_{QCD}^2)$. At small values of momentum transfer it should either vanish (if gluons decouple completely) or reach a constant

value [6, 7]. In the former case the gluon interaction would be screened at long range, which would most simply be achieved by taking $g(k^2)$ to have the form

$$g(k^2) = \frac{k^2}{\Lambda^2 + k^2} \frac{\log(\frac{4\Lambda^2}{\Lambda_{QCD}^2})}{\log(\frac{k^2 + 4\Lambda^2}{\Lambda_{QCD}^2})}, \quad (2.11)$$

where the parameter Λ , which represents the inverse of the screening length, is taken to be either Λ_{QCD} or Λ_χ . This in practice implies that the gluon attains a finite mass in the nonperturbative regime. For calculational convenience the expression (2.11) will be approximated here by the simpler form

$$g(k^2) = \frac{k^2}{\Lambda^2 + k^2} \frac{1}{1 + \frac{k^2}{\Lambda_F^2}}. \quad (2.12)$$

With an appropriately chosen value for Λ_F this form gives a fair approximation to (2.11) up to $k \simeq 5$ GeV. If the screening scale Λ is set to Λ_{QCD} (0.25 GeV [14]) and the effective coupling constant α_S in (2.9b) is taken to be $\alpha_S = 0.42$ and $\Lambda_F = 4.3$ GeV, the effective quark gluon coupling at the charmonium scale $k = 1.3$ GeV takes the value 0.37 and at the bottomonium scale $k = 4.1$ GeV the value 0.22 in agreement with the lattice calculation results values given in Ref. [15]. At the light constituent quark scale $k \simeq 0.34$ GeV the effective α_S value will then be $\alpha_S \simeq 0.27$, a value which is small enough that the spectroscopic problems associated with combinations of single gluon and pion exchange interactions are avoided [19]. If the screening scale Λ in (2.12) is set to $\Lambda_\chi \simeq 1$ GeV, the effective α_S value at the light constituent quark mass scale drops to only 0.04.

The fourth order terms in the quasipotential are the two-pion, two-gluon and the π -gluon exchange interactions:

$$U_{\pi\pi} = M_{\pi\pi} - K_\pi \tilde{g} K_\pi, \quad U_{GG} = M_{GG} - K_G \tilde{g} K_G, \quad (2.13a)$$

$$U_{\pi G} = M_{\pi G} - K_\pi \tilde{g} K_G - K_G \tilde{g} K_\pi. \quad (2.13b)$$

Here $M_{\pi\pi}$ and M_{GG} are the fourth order two-pion and two-gluon exchange quark-quark scattering amplitudes respectively and $M_{\pi G}$ is the fourth order π -gluon exchange amplitude. It is the latter one, and the corresponding quasipotential $U_{\pi G}$ that we shall consider here, because these have the same flavor dependence as the one-pion-exchange interaction K_π . The π -gluon exchange amplitude is illustrated by the Feynman diagrams in Fig. 1. The main part of the two-pion exchange interaction $K_{\pi\pi}$ is a central interaction with weak flavor dependence (cf. [12, 16]), which naturally is incorporated into the effective confining interaction between quarks. Similarly the two-gluon exchange interaction is flavor independent, and naturally incorporated into the confining interaction.

3. The π -Gluon Exchange Amplitude

The π -gluon exchange amplitude will have the same flavor dependence as the one-pion-exchange interaction (2.9a). Its spin structure may be expressed in terms of any set of linearly independent spin invariants. In order to avoid kinematic singularities, it proves convenient to employ the following set [9, 10]:

$$\begin{aligned} S_1 &= \gamma_5^{(1)} \gamma_5^{(2)} (\gamma^{(1)} \cdot \gamma^{(2)}), \quad S_2 = \{(\gamma^{(1)} \cdot N)(\gamma^{(2)} \cdot P), S_1\} \\ S_3 &= i[\gamma^{(1)} \cdot N + \gamma^{(2)} \cdot P]S_1, \quad S_4 = \gamma^{(1)} \cdot \gamma^{(2)} S_1, \quad S_5 = \gamma_5^{(1)} \gamma_5^{(2)}. \end{aligned} \quad (3.1)$$

Here the 4-vectors P and N have been defined in terms of the initial p_1, p_2 and final p'_1, p'_2 quark 4-momenta as

$$P = \frac{1}{2}(p_1 + p'_1), \quad N = \frac{1}{2}(p_2 + p'_2). \quad (3.2)$$

The relation between the invariants S_j and the more common Fermi invariants $SVTAP$ is given in Ref. [10].

The π -gluon exchange amplitude $M_{\pi G}$ may be expressed in terms of the spin invariants (3.1) as

$$M_{\pi G} = \sum_{j=1}^5 s_j(s, t, u) S_j \vec{\tau}^1 \cdot \vec{\tau}^2. \quad (3.3)$$

Here the variables s, t, u are defined as

$$s = -(p_1 + p_2)^2, \quad t = -(p'_1 - p_1)^2, \quad u = -(p'_1 - p_2)^2. \quad (3.4)$$

The scalar amplitudes s_j may be expressed in terms of dispersion relations in t as

$$s_j(s, t, u) = \frac{1}{\pi} \int_{t_0}^{\infty} dt' \frac{\sigma_j(t'; s, u)}{t' - t - i\epsilon}. \quad (3.5)$$

where $t_0 = (m_\pi + m_G)^2$. Here m_π is the pion mass and m_G is a fictitious gluon mass, which will be set to zero or to the screening parameter Λ (2.11) in the end. To derive the spectral functions $\sigma(t; s, u)$ we use the unitarity relation in the t -channel for the π -gluon intermediate states:

$$\sum_j \sigma_j S_j \vec{\tau}^1 \cdot \vec{\tau}^2 \simeq \frac{|\vec{q}|}{32\pi^2 \sqrt{t}} \int d\Omega_q T^\dagger T, \quad (3.6)$$

where T is the amplitude for gluoproduction of pions in the t -channel. The diagrams included in the unitarity integral are those shown in Fig. 1. Here the variable $|\vec{q}|$ is defined as

$$|\vec{q}| = \frac{1}{4t} [t^2 - 2t(m_\pi^2 + m_G^2) + (m_\pi^2 - m_G^2)^2]. \quad (3.7)$$

The ‘‘Born terms’’ in the pion gluoproduction amplitude T required in the unitarity integral (3.6) are readily obtained from the coupling Lagrangians (2.8): with exception of the color factor, the expressions are the same as those given in Ref. [10] for the $N\bar{N} \rightarrow \pi\omega$ amplitude. Note that the pion gluoproduction amplitude takes the same form for pseudovector and pseudoscalar π -quark coupling.

These “Born terms” in the pion gluoproduction amplitude then lead to the following expressions for the σ_j in (3.5), which may be obtained by suitable extension of the formalism in Ref. [10]:

$$\begin{aligned}
\sigma_1(s, t, u) &= m_q^2 \bar{N} [\theta_1(s, t) - \theta_1(u, t)], \\
\sigma_2(s, t, u) &= -\sigma_1(s, t, u)/2m_q^2, \\
\sigma_3(s, t, u) &= m_q \bar{N} [\theta_2(s, t) + \theta_2(u, t)], \\
\sigma_4(s, t, u) &= \bar{N} [\theta_4(s, t) + \theta_4(u, t)], \\
\sigma_5(s, t, u) &= \sigma_5^+(s, t, u) + \sigma_5^-(s, t, u), \\
\sigma_5^+(s, t, u) &= -\bar{N} [4\theta_4(s, t) + \frac{1}{4}(s - \bar{s})\theta_1(s, t) + (s \leftrightarrow u)], \\
\sigma_5^-(s, t, u) &= \bar{N} \{ I_0(s, t) [\frac{1}{4}(2s - 4m_q^2)(1 - \bar{\mu}^2/t)^2 + |\vec{q}|^2] \\
&\quad + (2m_q^2 - \frac{1}{4}t)\theta_1(s, t) - \frac{1}{2}(s - \bar{s})\theta_2(s, t) + \frac{1}{t}(s - \bar{s})\theta_4(s, t) - (s \leftrightarrow u) \}.
\end{aligned} \tag{3.8}$$

Here $\bar{N}$ is defined as

$$\bar{N} = -\frac{2f_{\pi qq}^2 \alpha_S}{3\pi} \frac{|\vec{q}|}{\sqrt{t}} \left(\frac{m_q}{m_\pi}\right)^2, \tag{3.9}$$

and $\bar{s} = 4m_q^2 - t - s$, with m_q being the quark mass.

The function I_0 in (3.8) is defined as

$$I_0(x, t) = \begin{cases} \frac{2\pi}{\sqrt{\xi^2 - \eta\zeta}} \log \frac{\xi + \zeta + \sqrt{\xi^2 - \eta\zeta}}{\xi + \zeta - \sqrt{\xi^2 - \eta\zeta}}, & \xi^2 - \eta\zeta > 0 \\ \frac{4\pi}{\xi + \zeta}, & \xi^2 - \eta\zeta = 0 \\ \frac{8\pi}{\sqrt{\zeta\eta - \xi^2}} \arctan \frac{2q^2 x}{\sqrt{\zeta\eta - \xi^2}}, & \xi^2 - \eta\zeta < 0. \end{cases} \tag{3.10}$$

Here the variables η , ξ , ζ , $\bar{\nu}$ and χ are defined as

$$\begin{aligned}
\eta &= 4|\vec{q}|^2 x, & \xi &= -2|\vec{q}|^2 x, \\
\zeta &= 4|\vec{q}|^2 \chi^2 + \frac{1}{4}(t - \bar{\nu}^2)^2, & \bar{\nu}^2 &= m_G^2 + m_\pi^2, \\
\chi^2 &= m_q^2 - \frac{1}{4}t.
\end{aligned} \tag{3.11}$$

The functions θ_j in (3.8) are defined as

$$\theta_1(x, t) = \frac{1}{x\bar{x}} \left\{ |\vec{q}|^2 (x - \bar{x}) I_0(x, t) + \frac{x + \bar{x}}{\chi^2} \left[\pi - \frac{1}{4} x' I(x') \right] - x' (2x - \bar{x}) R(x, t) \right\},$$

$$\begin{aligned}\theta_2(x, t) &= \frac{1}{x\bar{x}} \left\{ |\vec{q}|^2 (x + \bar{x}) I_0(x, t) + \frac{x - \bar{x}}{\chi^2} \left[\pi - \frac{1}{4} x' I(x') \right] - x' (2x + \bar{x}) R(x, t) \right\}, \\ \theta_4(x, t) &= |\vec{q}|^2 I_0(x, t) - x' R(x, t).\end{aligned}\tag{3.12}$$

Here the function $I(x')$ is defined as

$$I(x') = \begin{cases} \frac{2\pi}{|\vec{q}|\chi} \arctan \frac{2|\vec{q}|\chi}{x'}, & \chi^2 > 0 \\ \frac{\pi}{|\vec{q}|p} \log \left[\left(1 + \frac{2|\vec{q}|p}{x'} \right) / \left(1 - \frac{2|\vec{q}|p}{x'} \right) \right], & \chi^2 = -p^2 < 0, \end{cases}\tag{3.13}$$

with $\bar{x}$ and x' defined as:

$$\bar{x} = 4m_q^2 - t - x, \quad x' = \frac{1}{2}(t - \bar{\nu}^2).\tag{3.14}$$

Finally the function $R(x, t)$ in (3.12) is defined as

$$R(x, t) = \frac{1}{\bar{x}} [I(x') - x' I_0(x)].\tag{3.15}$$

For the definition of the π -gluon exchange potential the iterated single pion and gluon exchange interactions (2.9) have to be subtracted from these expressions (i.e. (3.5), (3.8)) for the π -gluon exchange amplitude.

4. The Iterated π -Gluon Exchange Interaction

The explicit form for the iterated π -gluon exchange interaction in (2.13b) is

$$\begin{aligned}J &= \frac{8\pi}{3} f_{\pi qq}^2 \alpha_s \frac{m_q^2}{m_\pi^2} \int \frac{d^2k}{(2\pi)^3} \frac{\vec{\tau}^1 \cdot \vec{\tau}^2}{E_k(k^2 - q^2 - i\epsilon)} \\ &\times \left\{ \frac{\gamma_5^{(1)} \gamma_5^{(2)} N(\vec{k}) \gamma^{(1)} \cdot \gamma^{(2)}}{[m_\pi^2 + (\vec{p}' - \vec{k})^2][m_G^2 + (\vec{p} - \vec{k})^2]} f[(\vec{p}' - \vec{k})^2] g[(\vec{p} - \vec{k})^2] \right. \\ &\left. + \frac{\gamma^{(1)} \cdot \gamma^{(2)} N(\vec{k}) \gamma_5^{(1)} \gamma_5^{(2)}}{[m_G^2 + (\vec{p}' - \vec{k})^2][m_\pi^2 + (\vec{p} - \vec{k})^2]} g[(\vec{p}' - \vec{k})^2] f[(\vec{p} - \vec{k})^2] \right\}.\end{aligned}\tag{4.1}$$

Here we have used the abbreviations

$$\begin{aligned}E_k &= \sqrt{k^2 + m_q^2}, \\ N(\vec{k}) &= (-\gamma^{(1)} \cdot \vec{k} - i\gamma_4^{(1)} E_k - im_q)(\gamma^{(2)} \cdot \vec{k} - i\gamma_4^{(2)} E_k - im_q).\end{aligned}\tag{4.2}$$

The spin-structure of the integral (4.1) may be expressed in terms of 5 spin amplitudes R_j defined in Ref. [9] as

$$\begin{aligned}R_1 &= \gamma_5^{(1)} \gamma_5^{(2)} (\gamma^{(1)} \cdot \gamma^{(2)}), \quad R_2 = \gamma_4^{(1)} \gamma_4^{(2)} R_1, \\ R_3 &= \{(\gamma_4^{(1)} + \gamma_4^{(2)}), R_1\}, \quad R_4 = \gamma^{(1)} \cdot \gamma^{(2)} R_1, \quad R_5 = \gamma_5^{(1)} \gamma_5^{(2)}.\end{aligned}\tag{4.3}$$

The integral (4.1) then takes the form

$$J = \sum_{i=1}^5 j_i(t, p^2) R_i \vec{\tau}^1 \cdot \vec{\tau}^2, \quad (4.4)$$

where the scalar amplitudes j_i may be expressed as dispersion integrals:

$$j_i(t, p^2) = \frac{1}{\pi} \int_{t_0}^{\infty} dt' \frac{\bar{\eta}_i(t', p^2)}{t' - t}. \quad (4.5)$$

Here the lower integration limit is $t_0 = (m_\pi + m_G)^2$.

The spectral functions $\bar{\eta}_i$ in these integrals have the explicit expressions

$$\begin{aligned} \bar{\eta}_1(t, p^2) &= \frac{4m_q^2\beta}{\kappa^4} \{I_1[|\vec{q}|^2(t - 4p^2) + 2\lambda^2] - 2\xi I_2 + 2I_3\}, \\ \bar{\eta}_2(t, p^2) &= \beta \left\{ E^2 I_1 \left[1 + \frac{1}{\kappa^2} \left(\frac{\bar{\mu}^4}{t} + 4p^2 - \frac{32|\vec{q}|^2 p^2}{\kappa^2} + \frac{8\lambda^2}{\kappa^2} \right) \right] \right. \\ &\quad \left. + I_2 \left[1 + \frac{4E^2}{\kappa^2} \left(1 - \frac{2\xi}{\kappa^2} \right) \right] + \frac{8E^2 I_3}{\kappa^4} - \frac{4E\pi}{\kappa^2} - \frac{2E(4p^2 + \bar{\nu}^2)}{\kappa^2} I_4 \right\}, \\ \bar{\eta}_3(t, p^2) &= \frac{m_q\beta}{2\kappa^2} \left\{ E I_1 \left(\bar{\nu}^2 - \frac{\bar{\mu}^4}{t} + \frac{32|\vec{q}|^2 p^2}{\kappa^2} - \frac{8\lambda^2}{\kappa^2} \right) \right. \\ &\quad \left. + 2E I_2 \left(\frac{4\xi}{\kappa^2} - 1 \right) - \frac{8E}{\kappa^2} I_3 + 2\pi + I_4(\bar{\nu}^2 - t) \right\}, \\ \bar{\eta}_4(t, p^2) &= \frac{\beta}{\kappa^2} \{ [4|\vec{q}|^2 p^2 - \lambda^2] I_1 + \xi I_2 - I_3 \}, \\ \bar{\eta}_5(t, p^2) &= \beta \left\{ I_1 \left[\delta \left(\frac{16|\vec{q}|^2 p^2 \gamma}{\kappa^2} - \frac{4\lambda^2 \gamma}{\kappa^2} + \frac{4p^2 \bar{\mu}^4}{\kappa^2 t^2} - \frac{4p^2}{\kappa^2} \right) - 2m_q^2 \left(\frac{4p^2 + \bar{\nu}^2}{\kappa^2} - \frac{\bar{\mu}^2}{t} \right) \right] \right. \\ &\quad \left. - \frac{I_2}{\kappa^2} [4m_q^2 + 4\delta(1 - \xi\gamma)] - \frac{4\delta\gamma}{\kappa^2} I_3 + \frac{8E\pi}{\kappa^2} + 4E I_4 \left[\frac{4p^2 + \bar{\nu}^2}{\kappa^2} - \frac{\bar{\mu}^2}{t} \right] \right\}. \end{aligned} \quad (4.6)$$

Here the notation is

$$\begin{aligned} \beta &= -\frac{2f_{\pi qq}^2 \alpha_S}{3\sqrt{t}} \left(\frac{m_q}{m_\pi} \right)^2, & \lambda^2 &= m_G^2 m_\pi^2, & \gamma &= \frac{2}{\kappa^2} + \frac{1}{t}, \\ \delta &= 2p^2 + m_q^2, & \xi &= t + p^2 - \bar{\nu}^2, & \bar{\mu}^2 &= m_G^2 - m_\pi^2, \\ \bar{\nu}^2 &= m_G^2 + m_\pi^2, & \kappa^2 &= 4p^2 + t, & E &= \sqrt{p^2 + m_q^2}. \end{aligned} \quad (4.7)$$

The functions I_k in these expressions are defined as

$$\begin{aligned}
I_1(t) &= \frac{1}{E} I_4 + \frac{2}{(p^2 - \sigma_-) \sqrt{\sigma_+ + m_q^2}} \left\{ \frac{\sigma_- + m_q^2}{E^2} \Pi(\gamma, \alpha) - F(\alpha) \right\}, \\
I_2(t) &= \frac{2}{\sqrt{\sigma_+ + m_q^2}} F(\alpha), \\
I_3(t) &= \frac{2}{\sqrt{\sigma_+ + m_q^2}} \{ (\sigma_+ + m_q^2) E(\alpha) - m_q^2 F(\alpha) \}, \\
I_4(t) &= \frac{\pi \sqrt{t}}{p} \left\{ \frac{\theta(t_+ - t)}{\sqrt{(t_+ - t)(t - t_-)}} + i \frac{\theta(t - t_+)}{\sqrt{(t - t_+)(t - t_-)}} \right\}.
\end{aligned} \tag{4.8}$$

Here $F(\alpha)$, $E(\alpha)$ and Π are complete elliptic integrals of the first, second and third kind, respectively, and the variables $\sigma_{\pm}$ are defined as

$$\sigma_{\pm} = \frac{1}{2} t' + p^2 - \frac{1}{2} \bar{\nu}^2 \pm \kappa q. \tag{4.9}$$

The arguments of the elliptic integrals are

$$\alpha = \frac{\sigma_+ - \sigma_-}{\sigma_+ + m_q^2}, \quad \gamma = \frac{p^2 - \sigma_-}{p^2 + m_q^2}. \tag{4.10}$$

In the expression for I_4 the variables $t_{\pm}$ are defined as

$$t_{\pm} = t_0 + \frac{\lambda^2}{2p^2} + 2\lambda \left\{ \pm \sqrt{1 + \frac{\bar{\nu}^2}{4p^2} + \frac{\lambda^2}{16p^4}} - 1 \right\}. \tag{4.11}$$

In order to construct the π -gluon exchange interaction (2.13b) the iterated π -gluon exchange interaction (4.1) has to be subtracted from the π -gluon exchange amplitude constructed in section 3 above. For that purpose the latter has to be expressed in terms of the spin operators R_j (4.3). This transformation has been given in Ref. [9]. In terms of these spin operators the π -gluon exchange amplitude (3.3) takes the form

$$M = \sum_{j=1}^5 r_j(s, t, u) R_j \vec{\tau}^1 \cdot \vec{\tau}^2. \tag{4.12}$$

The scalar amplitudes r_j are linear combinations of the amplitudes s_j in (3.3). These may also be written as dispersion integrals:

$$r_j(s, t, u) = \frac{1}{\pi} \int_{t_0}^{\infty} dt' \frac{\eta_j(t', s, u)}{t' - t}. \tag{4.13}$$

The spectral functions η_j here are the following linear combinations of the spectral functions σ_j in (3.8):

$$\begin{aligned}
\eta_1 &= \sigma_1 - 2m_q^2 \sigma_2 + 2m_q \sigma_3, & \eta_2 &= -8E^2 \sigma_2 + \sigma_4, \\
\eta_3 &= 2Em_q \sigma_2 - E \sigma_3, & \eta_4 &= \sigma_4, \\
\eta_5 &= (12E^2 - 2m_q^2) \sigma_2 + 2m_q \sigma_3 + \sigma_5.
\end{aligned} \tag{4.14}$$

Subtraction of the iterated π -gluon exchange interaction from the π -gluon exchange amplitude gives the final π -gluon exchange interaction in the form

$$U_{\pi G} = \sum_{j=1}^5 R_j \vec{\tau}^1 \cdot \vec{\tau}^2 \frac{1}{\pi} \int_{t_0}^{\infty} dt' \frac{\eta_j(s, t', u) - \bar{\eta}_j(t', s)}{t' - t}. \quad (4.15)$$

Here $s = 4m_q^2 + 4p^2$ in the argument of $\bar{\eta}_j$. Constructed in this way the π -gluon exchange interaction is real up to the pion production threshold, and only very weakly energy dependent.

5. The π -Gluon Exchange Potential

Constituent quarks confined within baryons and with masses in the 300-400 MeV range have large velocities and call for a Poincaré invariant treatment. The Blankenbecler-Sugar equation framework admits such a treatment, as demonstrated e.g. by the calculation of nucleon-nucleon scattering observables in Ref. [17]. In practice this calls for a helicity amplitude decomposition of the spin operators R_j (4.3) and solution of the resulting coupled linear integral equations. This approach, although straightforward, does not lead to very transparent results however. For a qualitative estimate of the numerical strength of the π -gluon exchange interaction, and of its importance in comparison to other interaction components, the standard adiabatic limit representation in terms of potential operators is most useful.

To lowest order in the nonlocal operator $(\vec{p}' + \vec{p})$ the potential may be expressed in terms of the potential operators:

$$\begin{aligned} \tilde{\Omega}_C &= 1, & \tilde{\Omega}_{LS} &= \frac{1}{2} i (\vec{\sigma}^1 + \vec{\sigma}^2) \cdot \vec{p}' \times \vec{p}, \\ \tilde{\Omega}_T &= \vec{\Delta}^2 (\vec{\sigma}^1 \cdot \vec{\sigma}^2) - 3 (\vec{\sigma}^1 \cdot \vec{\Delta}) (\vec{\sigma}^2 \cdot \vec{\Delta}), \\ \tilde{\Omega}_{SS} &= \vec{\sigma}^1 \cdot \vec{\sigma}^2, \end{aligned} \quad (5.1)$$

where $\vec{\Delta} = \vec{p}' - \vec{p}$. In the adiabatic limit the π -gluon exchange interaction then takes the following form, once the spin operators R_j are expanded in terms of the potential operators Ω_α :

$$v(\vec{p}', \vec{p}) = - \sum_{\alpha=1}^4 \sum_{j=1}^5 \vec{\tau}^1 \cdot \vec{\tau}^2 \tilde{\Omega}_\alpha X_{\alpha j} \frac{1}{\pi} \int_{t_0}^{\infty} dt' \frac{\eta_j - \bar{\eta}_j}{t' - t}. \quad (5.2)$$

Here $X_{\alpha j}$ is the linear transformation matrix between the spin operators R_j and the potential operators Ω_α :

$$X = \begin{bmatrix} 0 & 0 & 0 & \Delta^2/2m_q^2 & 0 \\ 1/2m_q^2 & -1/2m_q^2 & 0 & -5/2m_q^2 & 0 \\ -1/12m_q^2 & 1/6m_q^2 & 0 & 1/3m_q^2 & 1/12m_q^2 \\ -1 & -1 & -4 & -1 & -\Delta^2/12m_q^2 \end{bmatrix}. \quad (5.3)$$

For the visualization of the interaction it is useful to perform a Fourier transformation to configuration space, by which the potential takes the form

$$V(r) = \sum_{\alpha} v_{\alpha}(r) \Omega_{\alpha} \vec{\tau}^1 \cdot \vec{\tau}^2. \quad (5.4)$$

Here the spin operators Ω_{α} are defined in the conventional way as

$$\begin{aligned} \Omega_C &= 1, & \Omega_{LS} &= \frac{1}{2}(\vec{\sigma}^1 + \vec{\sigma}^2) \cdot \vec{L}, \\ \Omega_T &= 3(\vec{\sigma}^1 \cdot \hat{r})(\vec{\sigma}^2 \cdot \hat{r}) - \vec{\sigma}^1 \cdot \vec{\sigma}^2, \\ \Omega_{SS} &= \vec{\sigma}^1 \cdot \vec{\sigma}^2. \end{aligned} \quad (5.5)$$

For the potential functions $v_{\alpha}(r)$ the following explicit expressions obtain:

$$\begin{aligned} v_C(r) &= \frac{1}{4\pi^2} \int_{t_0}^{\infty} dt' \sqrt{t'} \eta_C(t') Y_0(r\sqrt{t'}), \\ v_{LS}(r) &= -\frac{1}{4\pi^2} \int_{t_0}^{\infty} dt' t' \sqrt{t'} \eta_{LS}(t') Y_0(r\sqrt{t'}) \left(1 + \frac{1}{r\sqrt{t'}}\right) / r\sqrt{t'}, \\ v_T(r) &= \frac{1}{4\pi^2} \int_{t_0}^{\infty} dt' t' \sqrt{t'} \eta_T(t') Y_0(r\sqrt{t'}) \left(1 + \frac{3}{r\sqrt{t'}} + \frac{3}{r^2 t'}\right), \\ v_{SS}(r) &= \frac{1}{4\pi^2} \int_{t_0}^{\infty} dt' \sqrt{t'} \eta_{SS}(t') Y_0(r\sqrt{t'}). \end{aligned} \quad (5.6)$$

Here Y_0 is the Yukawa function

$$Y_0(x) = e^{-x}/x. \quad (5.7)$$

The “mass spectrum” functions η_{α} are defined as

$$\eta_{\alpha}(t') = -\sum_{j=1}^5 X_{\alpha j} [\eta_j(t', s = 4m_q^2, u = 0) - \bar{\eta}_j(p^2 = 0, t')]. \quad (5.8)$$

Note that the integrals (5.6) are convergent even in the absence of a pion-quark vertex function. In that case the ultraviolet divergences in the dispersion integrals (3.5) and (4.5) may be isolated into delta functions by replacing the dispersion integrals by suitably subtracted versions.

6. Numerical Results

The controlling parameters for the π -gluon exchange interaction are the pion-quark coupling constant $f_{\pi qq}$, the coupling constant for gluons to constituent quarks α_S , the chiral restoration scale Λ_{χ} and the inverse screening length parameter Λ in the one-gluon exchange potential. Of these the pion-quark coupling constant and the chiral restoration scale parameters are probably the least contentious. The former may be related by standard quark model algebra to the pseudovector πNN coupling constant as $f_{\pi qq} = \frac{3}{5} f_{\pi NN}$ (see section 2 above). The value of the latter is $f_{\pi NN} \simeq 1$. For the chiral restoration scale the typical value is [18]

$$\Lambda_{\chi} \simeq 4\pi f_{\pi}, \quad (6.1)$$

where f_π is the pion decay constant. With $f_\pi=93$ MeV one obtains $\Lambda_\chi \simeq 1170$ MeV. We shall here use the value $\Lambda_\chi = 1000$ MeV.

The value of the coupling strength of gluons to constituent quarks α_S is not well established. We shall use $\alpha_S = 0.42$ in combination with the screening and fall-off factor (2.12) so as to achieve agreement with the lattice calculation results reported in Ref. [15] for momentum transfer values that correspond to the charmonium and bottomonium mass scales. As pointed out in section 2 this implies a very weak quark-gluon coupling for momenta that correspond to the light constituent quark mass scale (i.e. $\alpha_S \simeq 0.04 - 0.27$, depending on the choice of screening parameter Λ). Careful numerical studies [19] of the baryon spectrum with hyperfine interaction models that are linear combinations of the single gluon and pion exchange interactions [20, 21] indicate that α_S has to be less than about 0.35 if the problem of incorrect ordering of the lowest positive and negative parity states is to be avoided.

For the inverse screening length parameter Λ we shall use both the chiral restoration scale and the confinement scale Λ_{QCD} . As the latter is much smaller ($\simeq 250$ MeV [14]), the resulting π -gluon exchange interaction, with the exception of the spin-spin component, will be stronger for the latter value. The qualitative features will be the same however. There is obviously a possibility of balancing the values of α_S and the screening parameter against one another.

In Figs. 2 and 3 the central and spin-orbit components of the π -gluon exchange interaction are shown as functions of quark separation r , along with the corresponding Fourier transform of the components of the single gluon exchange interaction (note that the latter lack the flavor factor $\vec{\tau}_1 \cdot \vec{\tau}_2$). These components of the π -gluon exchange interaction have very short range (≤ 0.3 fm), and are weak. The most notable feature is the smallness of the spin-orbit component of the π -gluon exchange interaction. This is a phenomenologically desirable feature as the spin-orbit splitting of the P -shell of the baryon spectrum is very small. The smallness of the spin-orbit component is a consequence of the fact that the “mass spectrum” function η_{LS} in the spin-orbit potential $v_{LS}(r)$ (5.6) has a zero.

In Figs. 4 and 5 the tensor and spin-spin components of the π -gluon exchange interaction are given as functions of r . For comparison the corresponding components of the Fourier transform of the one-pion-exchange potential for quarks are also shown in Fig. 5. The most notable result here is that the strength of tensor component of the π -gluon exchange interaction is of the same order of magnitude as that of the one-pion-exchange interaction, although it has opposite sign. As a consequence the tensor component of the π -gluon exchange interaction tends to cancel that of the one-pion-exchange interaction. This cancellation of the main flavor dependent tensor interaction between the quarks provides a simple explanation for why the hyperfine splitting of the baryon spectrum is so well described by a simple pion exchange interaction, provided that its tensor component is dropped [2].

In the shortest range region the π -gluon tensor interaction is the predominant one, which is phenomenologically desirable, as the pion-exchange tensor by itself would split the spin-symmetric negative parity multiplet $N(1650) - N(1700) - N(1675)$ incorrectly [1].

The spin-spin component of the π -gluon exchange interaction has the same sign, and is of similar magnitude as the corresponding component of the one-pion-exchange interaction. This suggests that it is a significant component of the hyperfine interaction between quarks, even in the case of a very weak coupling between gluons and constituent quarks. This also contributes to the understanding of the phenomenological fact that the hyperfine splitting of the baryon spectrum can be explained by a strong attractive flavor dependent spin-spin interaction [1, 22].

The sign of the spin-spin component of the π -gluon exchange interaction changes at very short range. This sign change is a consequence of the weakening at high momentum transfer of the effective quark-gluon coupling constant (2.11), (2.12). As shown in Fig. 5 this feature is lost if the inverse screening length parameter Λ is set to the chiral restoration scale Λ_χ (1 GeV).

7. Discussion

The main result of the present investigation is that even in the case of very weak coupling of gluons to constituent quarks, the irreducible π -gluon exchange interaction is of significant magnitude. The reason for this is that even if the gluon exchange is screened for long wavelengths, the π -gluon exchange loop diagrams allow exchange of short wavelength gluons, along with long wavelength pions. The result indicates that the interaction between constituent quarks should be far more intricate than the conventional single gluon exchange and (less conventional for quarks) one-pion-exchange interactions. While the latter will be screened out at short range, the π -gluon exchange interaction is not restricted in that way.

The results suggest that the hyperfine interaction between light constituent quarks should be formed of a (weak) single gluon exchange component, a pion exchange component with conventional strength and an irreducible π -gluon exchange interaction. The single gluon exchange interaction is weak because the effective quark-gluon coupling is weak in the infrared limit [6, 7], and therefore the problem of a large gluonic spin-orbit interaction [3] is avoided. The pion exchange tensor interaction is in effect cancelled by the large tensor component of the π -gluon exchange interaction, and thus the incorrect (small) spin-orbit splitting of the low lying negative parity resonances is avoided. The π -gluon exchange and single pion exchange interactions combine to a strong attractive flavor dependent spin-spin interaction, which brings the low lying positive parity resonances below the lowest negative parity resonances in agreement with experiment.

There are a large number of other exchange mechanisms that may contribute significantly to the hyperfine interaction between quarks. Among these are vector meson exchange [1] and two-pion exchange, which definitely merit further study. The two-pion exchange interaction mainly leads to a flavor independent attractive interaction, which contributes to the strength of the effective confining interaction at short range. There will be a strong overlap between

that interaction and the vector meson exchange interactions.

The presence of an irreducible π –gluon exchange interaction is an immediate consequence if there exists a pion exchange and (however screened) gluon exchange interaction between quarks. It is also independent of the structure of the pion itself – whether it arises as a succession of instanton induced quark-antiquark interactions, according to a common view, or whether it has a simpler quark-antiquark structure.

There are several reasons to believe that the single gluon exchange interaction between constituent quarks should be very weak beyond the chiral restoration scale or at least beyond the confinement scale. Among these are the absence of gluonic degrees of freedom in cooled lattice calculations [4] and the very weak residual gluonic coupling in the valence-QCD approximation [5]. Phenomenological investigations of the behavior of the running coupling strength of quarks to gluons in the infrared limit also indicate that the coupling constant drops or freezes at a constant value [6, 7]. The (possible) disconnection of gluons from the constituent quarks at long range may be described phenomenologically in several ways. The most direct is the introduction of a suitable range dependence in the effective coupling constant α_S , which at very long range – i.e. beyond the confinement scale – weakens the coupling constant, which is known to increase with range in the short range region. In practice this would bring about the screening effect described here simply by the form (2.11).

The substantial size of the π –gluon tensor and spin-spin interaction components suggests that the interaction between constituent quarks in the end may prove to be as complex as the nucleon-nucleon interaction proved to be, and that it – at least partly – has to be constructed phenomenologically as is the case with the latter [23, 24].

Acknowledgments

D. O. R. thanks Professor S. J. Brodsky for helpful correspondence. This work has been supported in part by the Academy of Finland under contract 34081.

References

- [1] L. Ya. Glozman and D. O. Riska, Phys. Rept. **268**, 263 (1996).
- [2] L. Ya. Glozman et al., LANL hep-ph/9706507, unigraz-utp-25-06-97.
- [3] J. Carlson, J. B. Kogut and V. J. Pandharipande, Phys. Rev. D **28**, 2807 (1983).
- [4] M.-C. Chu et al., Phys. Rev. D **49**, 6039 (1994).
- [5] K.-F. Liu, private communication

- [6] A. C. Mattingly and P. M. Stevenson, Phys. Rev. D **49**, 437 (1994).
- [7] S. J. Brodsky et al., Phys. Rev. D **57**, 245 (1998).
- [8] R. Blankenbecler and R. Sugar, Phys. Rev. **142**, 1051 (1966).
- [9] D. O. Riska and Y. H. Chu, Nucl. Phys. **A235**, 499 (1974); **A251**, 530 (1975).
- [10] D. O. Riska, Nucl. Phys. **A274**, 349 (1976).
- [11] A. A. Logunov and A. N. Tavkhelidze, Nuovo Cim. **29**, 380 (1963).
- [12] M. H. Partovi and E. L. Lomon, Phys. Rev. D **2**, 1999 (1970).
- [13] F. Coester and D. O. Riska, Ann. Phys. **234**, 141 (1994).
- [14] J. F. Donoghue, E. Golowich and B. R. Holstein, Dynamics of the Standard Model, Cambridge University Press, Cambridge (1992).
- [15] C. T. H. Davies et al., Phys. Rev. D **56**, 2755 (1997).
- [16] M. Chemtob, J. W. Durso and D. O. Riska, Nucl. Phys. **B38**, 141 (1972).
- [17] A. D. Jackson, D. O. Riska and B. Verwest, Nucl. Phys. **A249**, 397 (1975).
- [18] A. Manohar and H. Georgi, Nucl. Phys. **B234**, 189 (1984).
- [19] L. Ya. Glozman et al., LANL nucl-th/9705011, unigraz-utp-05-05-97.
- [20] A. Valcarce et al., Phys. Lett. B **367**, 35 (1996).
- [21] Z. Dziembowski, M. Fabre de la Ripelle and G. A. Miller, Phys. Rev. C **53**, 2038 (1996).
- [22] F. Coester, K. Dannbom and D. O. Riska, LANL hep-ph/9711458.
- [23] M. Taketani, S. Nakamura and M. Sasaki, Prog. Theor. Phys. **6**, 581 (1951).
- [24] G. S. Bali, K. Schilling and A. Wachter, Phys. Rev. D **56**, 2566 (1997).

Figure Captions

Figure 1. Feynman diagram representation of the π -gluon exchange quark-quark scattering amplitude.





Figure 2. The central component of the π -gluon exchange potential as function of quark separation. The curve A is the result when the screening function $g(k^2)$ in the gluon exchange interaction is taken to be simply the fall-off factor $1/(1 + \frac{k^2}{\Lambda_F^2})$ and the curve B is the result when the full form (2.12) for the screening function is used with the screening

parameter Λ equal to Λ_{QCD} (250 MeV). The curve G(B) is the corresponding single gluon exchange interaction without the fall-off factor.

Figure 3. The spin-orbit component of the π -gluon exchange potential as function of quark separation. The curve A is the result when the screening function $g(k^2)$ in the gluon exchange interaction is taken to be simply the fall-off factor $1/(1 + \frac{k^2}{\Lambda_F^2})$ and the curve B is the result when the full form (2.12) for the screening function is used with the screening parameter Λ equal to Λ_{QCD} (250 MeV). The curve G(B) is the corresponding single gluon exchange interaction without the fall-off factor.

Figure 4. The tensor component of the π -gluon exchange potential as function of quark separation. The curve A is the result when the screening function $g(k^2)$ in the gluon exchange interaction is taken to be simply the fall-off factor $1/(1 + \frac{k^2}{\Lambda_F^2})$ and the curve B is the result when the full form (2.12) for the screening function is used with the screening parameter Λ equal to Λ_{QCD} (250 MeV). The curve pi is the corresponding one-pion exchange potential.

Figure 5. The spin-spin component of the π -gluon exchange potential as function of quark separation. The curve A is the result when the screening function $g(k^2)$ in the gluon exchange interaction is taken to be simply the fall-off factor $1/(1 + \frac{k^2}{\Lambda_F^2})$ and the curves B and C are the results when the full form (2.12) for the screening function is used with the screening parameter Λ equal to Λ_{QCD} (250 MeV) and Λ_χ (1 GeV), respectively. The curve pi is the corresponding one-pion exchange potential.






Fig. 3




Fig. 4




Fig. 5

